

# Large scale morphological changes in the Hapi region on Comet 67P/Churyumov-Gerasimenko

Björn J. R. Davidsson

Jet Propulsion Laboratory  
California Institute of Technology  
Pasadena (CA), USA

Seungwon Lee, Paul von Allmen, Pete Schloerb, Mark  
Hofstadter, Holger Sierks, Cesare Barbieri, Samuel  
Gulkis, Horst Uwe Keller, Detlef Koschny, Philippe Lamy,  
Hans Rickman, Rafa Rodrigo, the MIRO Team, the  
OSIRIS Team.




**Jet Propulsion Laboratory**  
California Institute of Technology

# Summary


- OSIRIS detected large-scale changes in the Hapi region
  - Shallow pits in the Hapi region formed in late Dec 2014, grew to 75 x 110 x 0.5 m in the next two months
  - Why did this happen?
- MIRO measured nucleus thermal emission at 1.59 mm and 0.53 mm
  - Temperature versus time and depth
  - Thermal inertia, ice abundance, extinction and scattering coefficients
  - We find a thermal inertia of 100-200 MKS
  - We find a drastic drop in water abundance prior to pit formation
    - In the range 11-22% ice by mass in Oct 2014
    - Perhaps as little as 2% ice by mass in Nov 2014
- Work in progress....

# Context



Credit: ESA/Rosetta/MPS for OSIRIS Team MPS/UPD/LAM/IAA/SSO/INTA/UPM/DASP/IDA and Davidsson *et al.* (2018, in preparation)

# A large shallow depression emerges




Dec 10, 2014. NAC 20 km:  $0.35 \text{ m px}^{-1}$ .




Jan 22, 2015. NAC 27km:  $0.49 \text{ m px}^{-1}$ .

# Accurate illumination conditions throughout orbit




WAC image on Feb 9, 2015,  
13:32:56.344 UTC

Credit: ESA/Rosetta/MPS for OSIRIS Team  
MPS/UPD/LAM/IAA/SSO/INTA/UPM/DASP/IDA



Synthetic image generated with the model of  
Davidsson & Rickman (2014, *Icarus* **243**, 58-77)  
Shape model SHAP5 version 1.5 (degraded)  
by Jorda *et al.* (2016, *Icarus*, **277**, 257-278)

# Accurate illumination conditions throughout orbit




Shape model with 50,000 facets

Direct solar illumination and shadowing by topography

Vis+IR self-illumination from surrounding terrain.

10° rotational steps throughout orbit.





1D heat conduction equation with upper boundary condition balancing illumination, thermal emission, heat conduction, ice sublimation.

Temperature  $T$  versus depth  $x$  and time  $t$ . Start with 90K during 1959 Jupiter encounter, integrate until present.


Credit: Davidsson



# Thermophysics and radiative transfer



Modeling the upper 3.4 meters.  
 $T=T(x)$  at different times during orbit.



$T(t)$  for different depths for  
 one comet day





Temperature profile function of

- \* Ice abundance
- \* Thermal inertia

Inserted into radiative transfer solver to calculate mm and smm radiances measured by MIRO, presented as *antenna temperature*.  
 Function of (per wavelength)





- \* Extinction coefficient
- \* Single-scattering albedo

# October 2014





# November 2014



# Summary

- The October 2014 MIRO measurements are consistent with 30%-50% water ice by volume, or 11-22% by mass (dust/ice mass ratio  $\mu = 3.5 - 8.1$ ).
- The November 2014 MIRO measurements are consistent with perhaps as little as 5% water ice by volume (~2% by mass).
- Throughout the period the thermal inertia is in the range 100-200 MKS
- Rapid loss of water ice (binding material, weight) just prior to pit formation may be related to their formation